

## **Colon Prep : GoLytely Extra Dose (2-Gallons)**

DOCTOR: Dr. Chua Dr. Cuschieri Dr. Ghorai Dr. Hawkins Dr. Loura Dr. Mu Dr. Schreiber Dr. Slosberg

Medical Group			
Calendar Dates			
5 DAYS PRIOR:	5 DAYS PRIOR:	1 DAY PRIOR:	PROCEDURE DATE:
General instructions	Medical Instructions	(Drink 1.5 gallon GoLytely)	(Drink 0.5 gallon GoLytely)
Last chance to cancel	From This Day On, Until The Procedure:	<b>7 AM</b> : Mix both GoLytely containers with	Early morning: take your <b>usual</b>
appointment or you will be	TURN PAPER OVER FOR LOW FIBER DIET	1 gallon of water each and store in	prescribed medication except for
charged a <b>NO SHOW FEE</b> of		refrigerator.	blood thinners, insulin or oral
\$250.		Entire Day: Clear liquid diet (no solids	hypoglycemic, unless told otherwise
	STOP:	foods!): clear fruit juice without pulp	by your doctor.
Check with insurance in	For days before procedure.	such as apple juice, white grape juice,	OK to continue clear liquid diet.
advance to see if you need		broth, Gatorade (No Gatorade Frost),	OK to continue clear liquid diet.
a referral or pre-approval	Stop taking oil supplements, oil capsules, iron pills,	popsicles, coffee or tea without cream.	SIX HOURS PRIOR TO YOUR CHECK
for your procedure.	as this makes it difficult to cleanse the bowel.	Small amount of Jell-O is OK. No milk,	IN TIME start drinking the remaining
Aa.a.a.alailalaana	If you are diabetic or if you're taking blood	red, blue, or purple colors.	GoLytely, one 8 oz. glass every 15 to
Arrange childcare, as children should not be	thinners and you did not receive specific	•	30 minutes until it is gone.
brought to the endoscopy	instructions regarding your medications, please call the office. Blood thinners include Plavix,	If you take diabetes medicine: Refer to	
center.	Effient, Warfarin (Coumadin), Pradaxa, or Xarelto,	the instruction sheet "Colonoscopy Patients with Diabetes".	DO NOT stop the prep even if you
center.	etc.	Patients with Diabetes .	think your stool is clear, as fecal
Secure a <b>driver</b> for the	etc.	<b>8 AM</b> : drink GoLytely, 8 oz. glass every 15	matter often comes out
procedure day, as it is	If you have implantable cardiology devices	to 30 minutes as tolerated. Finish 8	intermittently. A good prep should result in CLEAR yellowish watery
unsafe to drive for 12 hours	(defibrillator or pacemaker), notify the office as a	glasses or half gallon.	stool (not cloudy) that looks like
after the procedure.	device clearance is needed from your cardiologist.	<b>12 Noon</b> : drink the 2 <sup>nd</sup> half gallon of	clear urine.
	, , , , , , , , , , , , , , , , , , , ,	GoLytely, 8 oz. glass every 15 to 30	cicai diffic.
SUPPLIES:	You may take OTC MiraLAX one dose in a 8 oz.	minutes as tolerated until gone.	FOUR HOURS PRIOR TO YOUR
• 2 Gallons - GoLytely or a	glass of water up to 3 times per day as needed to	G	CHECK IN TIME STOP DRINKING ALL
generic equivalent	maintain daily soft BMs. You may also use your	<u>6 PM</u> : drink 2 <sup>nd</sup> gallon of GoLytely, 8 oz.	LIQUIDS.
(prescription)	usual laxatives as needed.	glass every 15 to 30 minutes as tolerated.	
MiraLAX (OTC), optional	If you have severe constipation, call office for	Finish 12 glasses and save the remaining	AGAIN, be sure to bring:
• Fleet enema (OTC),	advice (425) 259-3122.	4 glasses for the morning.	Medication list
optional		What to expect: It may take several	A driver      Dhote ID (Incurrence cond)
		hours before you have multiple bowel	Photo ID/Insurance card     Synglasses (Hearing side)
		movements. Expect lots of diarrhea.	<ul> <li>Eyeglasses/Hearing aids</li> </ul>
		*This is normal*	You <b>CANNOT</b> drink alcohol or drive
		NO ALCOHOL	for 12 hours after the test.

CPT Codes: Colonoscopy 45378-45385; ICD-10 codes: screening Z12.11; Hx Polyps Z86.010; Fx Polyps Z83.71; Fx Colon CA Z80.00; Other:\_\_\_\_\_\_

NEWLY REVISED 06/09/20