

Colon Prep: MiraLAX Extra Dose

Medical Group DOCTOR: Dr. Chua Dr. Cuschieri Dr. Ghorai Dr. Hawkins Dr. Loura Dr. Mu Dr. Schreiber Dr. Slosberg

Calendar Dates 5 DAYS PRIOR: 5 DAYS PRIOR: 1 DAY PRIOR: PROCEDURE DATE: (1^{st/}2 nd bottle of MiraLAX, 4 glasses/each) (3rd bottle of MiraLAX, 4 glasses) **General instructions Medical Instructions** Last chance to cancel From This Day On, Until The Procedure: Entire Day: Clear liquid diet only (no solids Early morning: take your usual prescribed appointment or you will be TURN PAPER OVER FOR LOW FIBER DIET medication except for blood thinners, foods!): clear fruit juice without pulp such as charged a **NO SHOW FEE** of apple juice, white grape juice, broth, insulin or oral hypoglycemic, unless told \$250. Gatorade (No Gatorade Frost), popsicles, otherwise by your doctor. STOP: coffee or tea without cream. Small amount Check with insurance in **For** days before procedure. of Jell-O is OK. No milk, red, blue, or purple OK to continue clear liquid diet. advance to see if you need a colors. referral or pre-approval for Stop taking oil supplements, oil capsules, iron **SIX HOURS PRIOR TO YOUR CHECK IN** your procedure. pills, as this makes it difficult to cleanse the If you take diabetes medicine: Refer to **TIME** Drink 3rd bottle MiraLAX, one 8 oz. the instruction sheet "Colonoscopy Patients bowel. glass every 20 minutes as tolerated. Finish Arrange childcare, as with Diabetes". 4 glasses until it is gone. children should not be If you are diabetic or if you're taking blood brought to the endoscopy 8 AM: Take 4 tablets of Dulcolax (Bisacodyl) thinners and you did not receive specific DO NOT stop the prep even if you think center. (OTC) instructions regarding your medications, your stool is clear, as fecal matter often Secure a **driver** for the please call the office. Blood thinners include comes out intermittently. A good prep 10 AM: Drink 1st bottle MiraLax/Gatorade. procedure day, as it is Plavix, Effient, Warfarin (Coumadin), Pradaxa, should result in CLEAR vellowish watery one 8 oz. glass every 20 minutes as unsafe to drive for 12 hours stool (not cloudy) that looks like clear or Xarelto, etc. tolerated. Finish 4 glasses until it is gone. after the procedure. urine. 6 PM: Drink 2nd bottle MiraLax/Gatorade, If you have implantable cardiology devices **SUPPLIES:** one 8 oz. glass every 20 minutes as **FOUR HOURS PRIOR TO YOUR CHECK IN** (defibrillator or pacemaker), notify the office • MiraLAX (OTC), 238gm tolerated. Finish 4 glasses until it is gone. TIME STOP DRINKING ALL LIQUIDS. as a device clearance is needed from your bottle/14 doses, buy cardiologist. **THREE bottles** What to expect: It may take several hours AGAIN, be sure to bring: • 32 oz. Bottle of Gatorade before you have multiple bowel movements. Medication list You may take OTC MiraLAX one dose in a 8 oz. (vellow or original Expect lots of diarrhea. A driver glass of water up to 3 times per day as needed only/no other colors), *This is normal* Photo ID/Insurance card to maintain daily soft BMs. You may also use **buy THREE bottles** NO ALCOHOL Eyeglasses/Hearing aids your usual laxatives as needed. • Dulcolax (Bisacodyl) 4 tablets (OTC) You **CANNOT** drink alcohol or drive for 12 Fleet enema (OTC), If you have severe constipation, call the office hours after the test. optional for advice (425) 259-3122.