



Diabetic Medication Instructions

Diabetic Medication	Morning Day Before Procedure	Night Before Procedure	Morning Day of Procedure
Oral: Traditional Tablets by mouth	Take usual dose if undergoing UPPER ENDOSCOPY	Do not take	Do not take (resume after procedure)
	Do not take if undergoing COLONOSCOPY		
Oral: New SGLT2 inhibitors <i>Jardiance (empagliflozin), Farxiga (dapagliflozin), Invokana (canagliflozin), Steglatro (ertugliflozin), Brenzavvy (bexagliflozin)</i>	This class of medication can cause ketoacidosis when patient has prolonged fasting. <ol style="list-style-type: none">1. If coming for upper endoscopy (EGD), hold evening dose the night before procedure and hold morning dose on day of procedure.2. If coming for a colonoscopy, stop taking 3 days prior to procedure date.		
Insulin- once daily (morning) <i>Levemir, Lantus, Trujoyo</i>	Take usual dose	N/A	Do not take (resume after procedure)
Insulin- once daily (evening) <i>Levemir, Lantus, Trujoyo</i>	N/A	Take 1/2 dose	Do not take (resume after procedure)
Insulin- twice daily <i>Humulin-N, Novolin-N</i>	Take half of your usual dose	Do not take	Do not take (resume after procedure)
Insulin Pump	Confirm dosage adjustment with your diabetes clinic		
Short-acting insulin <i>(Humalog, Novolog, Apidra)</i>	Take the usual amount of regular insulin for blood sugars >220mg/dL		
GLP-1 agonists <i>(Byetta, Saxenda, Victoza, Adlyixin, Rybelsus, Trulicity, Wegovy, Ozempic, Bydureon, BCise, Mounjaro)</i>	<ol style="list-style-type: none">1. If you're receiving a GLP-1 agonist once daily, stop using it one day before the GI procedure(s).2. If you're receiving a GLP-1 agonist once weekly, stop using it for 7 days before the GI procedure(s).3. Note: When the GLP-1 agonist is withheld, your blood sugar level may go up slightly. Monitor your blood sugar level closely and make adjustments of your other diabetic medications as needed.		